

# Broadcast domination and multipacking in graphs and digraphs

**Florent Foucaud**

(Université de Bordeaux, France)

joint works with:

**Laurent Beaudou**

(Higher School of Economics, Moscow, Russia)

**Richard C. Brewster**

(Thompson Rivers University, Kamloops, BC, Canada)

**Benjamin Gras**

(Université d'Orléans, France)

**Anthony Perez**

(Université d'Orléans, France)


**Florian Sikora**

(Université Paris-Dauphine, France)

Covering: cover the vertices of a graph using as few structures as possible

*Example: dominating set:* covering using 1-balls

→ domination number  $\gamma(G)$



**Covering:** cover the vertices of a graph using as few structures as possible

*Example: dominating set:* covering using 1-balls


→ domination number  $\gamma(G)$



**Packing:** pack as many structures as possible without interference

*Example: dist. 3-independent set / 2-packing:* packing 1-balls without overlap

→ 2-packing number  $\rho_2(G)$



**Covering:** cover the vertices of a graph using as few structures as possible

*Example:* dominating set: covering using 1-balls


→ domination number  $\gamma(G)$



**Packing:** pack as many structures as possible without interference

*Example:* dist. 3-independent set / 2-packing: packing 1-balls without overlap

→ 2-packing number  $\rho_2(G)$



These problems are **dual** (in the sense of LP) and  $\rho_2(G) \leq \gamma(G)$ .


**Definition** - Dominating broadcast of graph  $G$  (Erwin, 2001)

A function  $f : V(G) \rightarrow \mathbb{N}$  s.t. for every  $v \in V(G)$ , there exists  $x \in V(G)$  with

- $f(x) > 0$  and
- $f(x) \geq d_G(x, v)$ .

The cost of  $f$  is  $\sum_{v \in V(G)} f(v)$ .

Broadcast number  $\gamma_b(G)$ : smallest cost of a dominating broadcast of  $G$ .




## Definition - Dominating broadcast of graph $G$ (Erwin, 2001)

A function  $f : V(G) \rightarrow \mathbb{N}$  s.t. for every  $v \in V(G)$ , there exists  $x \in V(G)$  with

- $f(x) > 0$  and
- $f(x) \geq d_G(x, v)$ .

The cost of  $f$  is  $\sum_{v \in V(G)} f(v)$ .

Broadcast number  $\gamma_b(G)$ : smallest cost of a dominating broadcast of  $G$ .




## Definition - Dominating broadcast of graph $G$ (Erwin, 2001)

A function  $f : V(G) \rightarrow \mathbb{N}$  s.t. for every  $v \in V(G)$ , there exists  $x \in V(G)$  with

- $f(x) > 0$  and
- $f(x) \geq d_G(x, v)$ .

The cost of  $f$  is  $\sum_{v \in V(G)} f(v)$ .

Broadcast number  $\gamma_b(G)$ : smallest cost of a dominating broadcast of  $G$ .




## Definition - Dominating broadcast of graph $G$ (Erwin, 2001)

A function  $f : V(G) \rightarrow \mathbb{N}$  s.t. for every  $v \in V(G)$ , there exists  $x \in V(G)$  with

- $f(x) > 0$  and
- $f(x) \geq d_G(x, v)$ .

The cost of  $f$  is  $\sum_{v \in V(G)} f(v)$ .

Broadcast number  $\gamma_b(G)$ : smallest cost of a dominating broadcast of  $G$ .






## Definition - Dominating broadcast of graph $G$ (Erwin, 2001)

A function  $f : V(G) \rightarrow \mathbb{N}$  s.t. for every  $v \in V(G)$ , there exists  $x \in V(G)$  with

- $f(x) > 0$  and
- $f(x) \geq d_G(x, v)$ .

The cost of  $f$  is  $\sum_{v \in V(G)} f(v)$ .

Broadcast number  $\gamma_b(G)$ : smallest cost of a dominating broadcast of  $G$ .



## Theorem (Heggernes-Lokshtanov, 2006)

We can find a minimum-cost dominating broadcast in polynomial time  $O(n^6)$ .


Proof idea:

- find an **efficient** dominating broadcast (Erwin, 2001)
- The structure of covering balls is a path or a cycle
- Dynamic programming on this structure

**Definition** - Multipacking of graph  $G$  (Brewster-Mynhardt-Teshima, 2014)

A set  $S$  of vertices s.t. for every  $v \in V(G)$  and every  $d \in \mathbb{N}$ , the  $d$ -ball  $B_d(v)$  contains at most  $d$  vertices of  $S$ .


Multipacking number  $mp(G)$ : largest size of a multipacking of  $G$ .



**Definition** - Multipacking of graph  $G$  (Brewster-Mynhardt-Teshima, 2014)

A set  $S$  of vertices s.t. for every  $v \in V(G)$  and every  $d \in \mathbb{N}$ , the  $d$ -ball  $B_d(v)$  contains at most  $d$  vertices of  $S$ .


Multipacking number  $mp(G)$ : largest size of a multipacking of  $G$ .



**Definition** - Multipacking of graph  $G$  (Brewster-Mynhardt-Teshima, 2014)

A set  $S$  of vertices s.t. for every  $v \in V(G)$  and every  $d \in \mathbb{N}$ , the  $d$ -ball  $B_d(v)$  contains at most  $d$  vertices of  $S$ .

**Multipacking number**  $mp(G)$ : largest size of a multipacking of  $G$ .



# Bounds for undirected graphs

The two problems are **dual** (in the sense of LP).

## Proposition

For every graph  $G$ , we have  $mp(G) \leq \gamma_b(G)$ .

Equality holds for:

- **trees** (Mynhardt-Teshima, 2017)
- more generally, **strongly chordal graphs** (Brewster-MacGillivray-Yang, 2019)
- **square grids** (Beaudou-Brewster, 2018)

**Proposition** (Erwin, 2001 + Hartnell-Mynhardt, 2014)


For any graph  $G$ ,  $\left\lceil \frac{\text{diam}(G)+1}{3} \right\rceil \leq mp(G) \leq \gamma_b(G) \leq \text{rad}(G) \leq \text{diam}(G)$ .



**Proposition** (Erwin, 2001 + Hartnell-Mynhardt, 2014)

For any graph  $G$ ,  $\left\lceil \frac{\text{diam}(G)+1}{3} \right\rceil \leq mp(G) \leq \gamma_b(G) \leq \text{rad}(G) \leq \text{diam}(G)$ .


$\gamma_b(G) \leq \text{rad}(G)$ : consider a **radial vertex**  $v$ . Set  $f(v) = \text{rad}(G)$ .



**Proposition** (Erwin, 2001 + Hartnell-Mynhardt, 2014)

For any graph  $G$ ,  $\left\lceil \frac{\text{diam}(G)+1}{3} \right\rceil \leq mp(G) \leq \gamma_b(G) \leq \text{rad}(G) \leq \text{diam}(G)$ .

$\left\lceil \frac{\text{diam}(G)+1}{3} \right\rceil \leq mp(G)$ : consider a **diametral path**  $P$ , select every third vertex.



**Proposition** (Erwin, 2001 + Hartnell-Mynhardt, 2014)

For any graph  $G$ ,  $\left\lceil \frac{\text{diam}(G)+1}{3} \right\rceil \leq mp(G) \leq \gamma_b(G) \leq \text{rad}(G) \leq \text{diam}(G)$ .

**Corollary**

For any graph  $G$ , we have  $\gamma_b(G) < 3mp(G)$ , hence  $\frac{\gamma_b(G)}{mp(G)} < 3$ .

**Question** (Hartnell-Mynhardt, 2014)

What is the largest possible ratio  $\frac{\gamma_b(G)}{mp(G)}$ ?

**Theorem** (Beaudou, Brewster, F., 2018)

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G) + 3$ , hence  $\frac{\gamma_b(G)}{mp(G)} \leq 2 + \epsilon$ .

**Theorem** (Beaudou, Brewster, F., 2018)


For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G) + 3$ , hence  $\frac{\gamma_b(G)}{mp(G)} \leq 2 + \epsilon$ .

**Lemma****Proof sketch**

Let  $u, v, x, y$  be 4 vertices with:

- $d(u, v) = 6k$
- $d(x, u) = d(x, v) = 3k$
- $d(x, y) = 3k + 3\ell$ .

Then,  $mp(G) \geq 2k + \ell$ .



**Theorem** (Beaudou, Brewster, F., 2018)

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G) + 3$ , hence  $\frac{\gamma_b(G)}{mp(G)} \leq 2 + \epsilon$ .

**Lemma****Proof sketch**

Let  $u, v, x, y$  be 4 vertices with:

- $d(u, v) = 6k$
- $d(x, u) = d(x, v) = 3k$
- $d(x, y) = 3k + 3\ell$ .

Then,  $mp(G) \geq 2k + \ell$ .

Let  $diam(G) = 6k + i$  and  $rad(G) = 3k + 3\ell + j$

( $0 \leq i < 6$  and  $0 \leq j < 3$ )

Apply the lemma with  $x$ , a vertex of eccentricity  $rad(G)$ .

$$\begin{aligned}
 mp(G) &\geq 2k + \ell \\
 &\geq \frac{diam(G)}{3} + \frac{rad(G)}{3} - \frac{diam(G)}{6} - c \\
 &\geq \frac{rad(G)}{2} - c \\
 &\geq \frac{\gamma_b(G)}{2} - c
 \end{aligned}$$

□

## Theorem (Beaudou, Brewster, F., 2018)

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G) + 3$ , hence  $\frac{\gamma_b(G)}{mp(G)} \leq 2 + \epsilon$ .

## Conjecture

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Theorem (Beaudou, Brewster, F., 2018)

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G) + 3$ , hence  $\frac{\gamma_b(G)}{mp(G)} \leq 2 + \epsilon$ .

## Conjecture

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Theorem (Beaudou, Brewster, F., 2018)

The conjecture is true when  $mp(G) \leq 4$ .



**Theorem** (Beaudou, Brewster, F., 2018)

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G) + 3$ , hence  $\frac{\gamma_b(G)}{mp(G)} \leq 2 + \epsilon$ .




**Conjecture**

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

**Theorem** (Beaudou, Brewster, F., 2018)

The conjecture is true when  $mp(G) \leq 4$ .

Conjecture would be tight — infinitely many graphs  $G$  s.t.  $\gamma_b(G) = 2mp(G)$ :



$$mp(G) = 2 \text{ and } \gamma_b(G) = 4$$

## Conjecture

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Question

What happens for **connected** graphs?


## Conjecture

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Question

What happens for **connected** graphs?

Closest known **connected** family:  $\gamma_b(G) = \frac{4}{3}mp(G)$  (Hartnell-Mynhardt, 2014)



Here:  $mp(G) = 3$  and  $\gamma_b(G) = 4$ .


## Conjecture

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Question

What happens for **connected** graphs?

Closest known **connected** family:  $\gamma_b(G) = \frac{4}{3}mp(G)$  (Hartnell-Mynhardt, 2014)



Here:  $mp(G) = 3$  and  $\gamma_b(G) = 4$ .


## Conjecture

For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Question

What happens for **connected** graphs?

Closest known **connected** family:  $\gamma_b(G) = \frac{4}{3}mp(G)$  (Hartnell-Mynhardt, 2014)



Here:  $mp(G) = 3$  and  $\gamma_b(G) = 4$ .

## Conjecture


For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Question

What happens for **connected** graphs?

## Proposition

There is a connected graph  $G$  of order 24 with  $mp(G) = 3$  and  $\gamma_b(G) = 5$ .



## Conjecture


For any graph  $G$ , we have  $\gamma_b(G) \leq 2mp(G)$ .

## Question

What happens for **connected** graphs?

## Proposition

There is a connected graph  $G$  of order 24 with  $mp(G) = 3$  and  $\gamma_b(G) = 5$ .





## Question

Is there a **connected** graph  $G$  with  $mp(G) = 3$  and  $\gamma_b(G) = 6$ ?


# Complexity & Algorithms for directed graphs







Note: an **undirected** graph can be seen as a **symmetric** directed graph!



Broadcast domination for directed graphs:

A vertex  $v$  with  $f(v) = r$  broadcasts to all vertices at **directed distance** up to  $r$ .

## BROADCAST DOM

Input: A (directed) graph  $G$ , an integer  $k$ .

Question: Does  $G$  have a dominating broadcast of cost at most  $k$ ?

**Theorem** (Heggernes-Lokshtanov, 2006)

BROADCAST DOM can be solved in polynomial time  $O(n^6)$  for **undirected graphs**.

## BROADCAST DOM

Input: A (directed) graph  $G$ , an integer  $k$ .

Question: Does  $G$  have a dominating broadcast of cost at most  $k$ ?



**Theorem** (Heggernes-Lokshtanov, 2006)

BROADCAST DOM can be solved in polynomial time  $O(n^6)$  for **undirected graphs**.

**Theorem** (F., Gras, Perez, Sikora, 2019)

BROADCAST DOM is NP-hard and  $W[2]$ -hard: likely no algorithm of the form  $f(k)poly(n)$ , for any computable function  $f$ .

Proof: Reductions from SET COVER.



## BROADCAST DOM

Input: A (directed) graph  $G$ , an integer  $k$ .

Question: Does  $G$  have a dominating broadcast of cost at most  $k$ ?

**Theorem** (F., Gras, Perez, Sikora, 2019)

BROADCAST DOM is NP-hard and  $W[2]$ -hard: likely no algorithm of the form  $f(k)poly(n)$ , for any computable function  $f$ .

## BROADCAST DOM

Input: A (directed) graph  $G$ , an integer  $k$ .

Question: Does  $G$  have a dominating broadcast of cost at most  $k$ ?

**Theorem** (F., Gras, Perez, Sikora, 2019)

BROADCAST DOM is NP-hard and  $W[2]$ -hard: likely no algorithm of the form  $f(k)\text{poly}(n)$ , for any computable function  $f$ .

**Theorem** (F., Gras, Perez, Sikora, 2019)

There is an  $O(c^k n)$ -time algorithm for BROADCAST DOM for **directed acyclic graphs**.

**Theorem** (F., Gras, Perez, Sikora, 2019)

There is a linear-time algorithm for BROADCAST DOM on **single-source layered directed graphs**.

### MULTIPACKING

Input: A (directed) graph  $G$ , an integer  $k$ .

Question: Does  $G$  have a multipacking of size at least  $k$ ?

(Note: OPEN for undirected graphs.)



## MULTIPACKING

Input: A (directed) graph  $G$ , an integer  $k$ .


Question: Does  $G$  have a multipacking of size at least  $k$ ?

(Note: OPEN for undirected graphs.)

**Theorem** (F., Gras, Perez, Sikora, 2019)

MULTIPACKING is NP-hard and  $W[1]$ -hard: likely no algorithm of the form  $f(k)poly(n)$ , for any computable function  $f$ .

Proof: Reduction from INDEPENDENT SET.



**Theorem** (F., Gras, Perez, Sikora, 2019)

MULTIPACKING is NP-hard and  $W[1]$ -hard: likely no algorithm of the form  $f(k)poly(n)$ , for any computable function  $f$ .

**Theorem** (F., Gras, Perez, Sikora, 2019)

MULTIPACKING is NP-hard and  $W[1]$ -hard: likely no algorithm of the form  $f(k)\text{poly}(n)$ , for any computable function  $f$ .

**Theorem** (F., Gras, Perez, Sikora, 2019)

There is a linear-time algorithm for MULTIPACKING on [single-source layered directed graphs](#).

Bounds:

- Is the conjecture true that for any **undirected** graph  $G$ ,  $\gamma_b(G) \leq 2mp(G)$ ?
- What is a tight bound for **connected undirected** graphs?  $\gamma_b(G) \leq \frac{4}{3}mp(G)$ ?
- What about **directed graphs**?

Complexity:

- Is MULTIPACKING NP-hard on **undirected** graphs?
- Complexity of both problems on **oriented trees**?

Bounds:

- Is the conjecture true that for any **undirected** graph  $G$ ,  $\gamma_b(G) \leq 2mp(G)$ ?
- What is a tight bound for **connected undirected** graphs?  $\gamma_b(G) \leq \frac{4}{3}mp(G)$ ?
- What about **directed graphs**?

Complexity:

- Is MULTIPACKING NP-hard on **undirected** graphs?
- Complexity of both problems on **oriented trees**?

# Thanks!